arXiv:hep-ph/9605440v3 9 Oct 1996

LTH 372
hep-ph/9605440
Revised Version

The three-loop SSM β-functions

P.M. Ferreira, I. Jack and D.R.T. Jones

Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, U.K.

We present the supersymmetric standard model three-loop β-functions for gauge and Yukawa couplings and consider the effect of three-loop corrections on the standard running coupling analyses.

May 1996

The unification (or near–unification) of the gauge couplings at $M_G \approx 10^{16}$ GeV has catalysed intensive study of the supersymmetric standard model (SSM). The evolution of the Yukawa couplings from M_Z to M_G is also of interest. For small $\tan\beta$, the t-quark Yukawa coupling y_t exhibits quasi–infrared fixed point (QFP) behaviour [1], and the corresponding form of $y_t(\mu)$ is favourable for b/τ unification at M_G. For large $\tan\beta$, trinification ($y_t(M_G) = y_b(M_G) = y_\tau(M_G)$) is possible[2]. Contemporary analyses generally employ the two-loop gauge and Yukawa β-functions, and apply one loop corrections in the low energy theory. In general the change in low energy predictions resulting from use of two-loop rather than one-loop β-functions is quite small; it appears that perturbation theory is reliable. In this paper we take the first step beyond these calculations by presenting the three loop β-functions for the dimensionless couplings. We deduce the three-loop Yukawa β-functions from the recent dimensional reduction (DRED) calculation of the three loop anomalous dimension of the chiral supermultiplet in a general $N = 1$ theory [3]. We also need the three-loop gauge β-functions. In Ref. [4] it is shown by explicit calculation that in the abelian case the DRED result for $\beta_g^{(3)}$ differs from that obtained from the exact NSVZ formula of Ref.[5] by a simple coupling constant redefinition, and the corresponding redefinition for the non-abelian case is inferred. From Ref. [4] we are thus able to obtain the DRED results for the three-loop gauge β-functions. In fact the effect of using these rather than the results of Ref. [5] is very small in the examples we present here.

We run gauge and Yukawa couplings between M_Z and M_G and compare the results with the corresponding calculations at one and two loops. In general the three-loop effects are small and opposite in sign to the two-loop ones; certainly not more significant than one- and two-loop radiative corrections at M_Z (some of which are not yet calculated). We do not, therefore, perform detailed phenomenological analysis. The relevant part of the SSM superpotential is:

$$W = H_2 t Y_t Q + H_1 b Y_b Q + H_1 \tau Y_\tau L \tag{1}$$

where Y_t, Y_b, Y_τ are $n_g \times n_g$ Yukawa matrices, and we define

$$T = Y_t^\dagger Y_t \, , B = Y_b^\dagger Y_b \, , E = Y_\tau^\dagger Y_\tau \, , \tilde{T} = Y_t \, Y_t^\dagger , \tilde{B} = Y_b \, Y_b^\dagger , \tilde{E} = Y_\tau \, Y_\tau^\dagger . \tag{2}$$

The $SU_3 \otimes \; SU_2 \otimes \; U_1$ gauge β-functions are as follows:

$$\beta_{g_i} = (16\pi^2)^{-1} b_i g_i^3 + (16\pi^2)^{-2} g_i^3 \left(\sum_j b_{ij} g_j^2 - a_i \right) + (16\pi^2)^{-3} \beta_{g_i}^{(3)} + \cdots \tag{3}$$

where

$$\begin{aligned}
&b_1 = 2n_g + \tfrac{3}{5}, \quad b_2 = 2n_g - 5, \quad b_3 = 2n_g - 9 \\
&a_1 = \tfrac{26}{5}\mathrm{tr}T + \tfrac{14}{5}\mathrm{tr}B + \tfrac{18}{5}\mathrm{tr}E, \quad a_2 = 6\mathrm{tr}T + 6\mathrm{tr}B + 2\mathrm{tr}E, \quad a_3 = 4\mathrm{tr}T + 4\mathrm{tr}B
\end{aligned} \tag{4}$$

and

$$b_{ij} = \begin{pmatrix} \frac{38}{15}n_g + \frac{9}{25} & \frac{6}{5}n_g + \frac{9}{5} & \frac{88}{15}n_g \\ \frac{2}{5}n_g + \frac{3}{5} & 14n_g - 17 & 8n_g \\ \frac{11}{15}n_g & 3n_g & \frac{68}{3}n_g - 54 \end{pmatrix}. \tag{5}$$

The three loop terms are[4]:

$$\begin{aligned}
\beta_{g_1}^{(3)} =& g_1^3\Big[\tfrac{84}{5}\mathrm{tr}T^2 + 18(\mathrm{tr}T)^2 + \tfrac{54}{5}\mathrm{tr}B^2 + \tfrac{36}{5}(\mathrm{tr}B)^2 + \tfrac{58}{5}\mathrm{tr}TB + \tfrac{54}{5}\mathrm{tr}E^2 + \tfrac{24}{5}(\mathrm{tr}E)^2 \\
&+ \tfrac{84}{5}\mathrm{tr}E\mathrm{tr}B - \left(\tfrac{169}{75}g_1^2 + \tfrac{87}{5}g_2^2 + \tfrac{352}{15}g_3^2\right)\mathrm{tr}T - \left(\tfrac{49}{75}g_1^2 + \tfrac{33}{5}g_2^2 + \tfrac{256}{15}g_3^2\right)\mathrm{tr}B \\
&- (\tfrac{81}{25}g_1^2 + \tfrac{63}{5}g_2^2)\mathrm{tr}E - (\tfrac{88}{5}n_g^2 - \tfrac{572}{9}n_g)g_3^4 - (\tfrac{18}{5}n_g^2 - \tfrac{9}{5}n_g - \tfrac{54}{5})g_2^4 \\
&- (\tfrac{38}{5}n_g^2 + \tfrac{1261}{225}n_g + \tfrac{54}{125})g_1^4 - \tfrac{1096}{225}n_g g_3^2 g_1^2 - \tfrac{8}{5}n_g g_2^2 g_3^2 - (\tfrac{27}{25} + \tfrac{14}{25}n_g)g_1^2 g_2^2\Big],
\end{aligned} \tag{6a}$$

$$\begin{aligned}
\beta_{g_2}^{(3)} =& g_2^3\Big[24\mathrm{tr}T^2 + 18(\mathrm{tr}T)^2 + 24\mathrm{tr}B^2 + 18(\mathrm{tr}B)^2 + 12\mathrm{tr}BT + 12\mathrm{tr}B\mathrm{tr}E \\
&+ 8\mathrm{tr}E^2 + 2(\mathrm{tr}E)^2 - \left(32g_3^2 + 33g_2^2 + \tfrac{29}{5}g_1^2\right)\mathrm{tr}T - \left(32g_3^2 + 33g_2^2 + \tfrac{11}{5}g_1^2\right)\mathrm{tr}B \\
&- \left(11g_2^2 + \tfrac{21}{5}g_1^2\right)\mathrm{tr}E - (24n_g^2 - \tfrac{260}{3}n_g)g_3^4 - (26n_g^2 - 123n_g + 100)g_2^4 \\
&- (\tfrac{6}{5}n_g^2 + \tfrac{169}{75}n_g + \tfrac{18}{25})g_1^4 + (\tfrac{2}{5}n_g + \tfrac{3}{5})g_1^2 g_2^2 + 8n_g g_2^2 g_3^2 - \tfrac{8}{15}n_g g_1^2 g_3^2\Big],
\end{aligned} \tag{6b}$$

$$\begin{aligned}
\beta_{g_3}^{(3)} =& g_3^3\Big[18(\mathrm{tr}T)^2 + 12\mathrm{tr}T^2 + 8\mathrm{tr}BT + 12(\mathrm{tr}B)^2 + 18\mathrm{tr}B^2 + 6\mathrm{tr}E\mathrm{tr}B \\
&- (\tfrac{104}{3}g_3^2 + 12g_2^2)(\mathrm{tr}T + \mathrm{tr}B) - g_1^2(\tfrac{44}{15}\mathrm{tr}T + \tfrac{32}{15}\mathrm{tr}B) - (44n_g^2 - \tfrac{3236}{9}n_g + 567)g_3^4 \\
&+ 2n_g g_3^2 g_2^2 + \tfrac{22}{45}n_g g_3^2 g_1^2 - (\tfrac{11}{5}n_g^2 + \tfrac{217}{225}n_g)g_1^4 - (9n_g^2 - 18n_g)g_2^4 - \tfrac{1}{5}n_g g_1^2 g_2^2\Big].
\end{aligned} \tag{6c}$$

For the anomalous dimensions of the chiral superfields we have at one loop:

$$\begin{aligned}
16\pi^2\gamma_t^{(1)} &= 2\tilde{T} - \tfrac{8}{3}g_3^2 - \tfrac{8}{15}g_1^2, \\
16\pi^2\gamma_b^{(1)} &= 2\tilde{B} - \tfrac{8}{3}g_3^2 - \tfrac{2}{15}g_1^2, \\
16\pi^2\gamma_Q^{(1)} &= B + T - \tfrac{8}{3}g_3^2 - \tfrac{3}{2}g_2^2 - \tfrac{1}{30}g_1^2, \\
16\pi^2\gamma_{H_2}^{(1)} &= 3\mathrm{tr}T - \tfrac{3}{2}g_2^2 - \tfrac{3}{10}g_1^2, \\
16\pi^2\gamma_{H_1}^{(1)} &= \mathrm{tr}E + 3\mathrm{tr}B - \tfrac{3}{2}g_2^2 - \tfrac{3}{10}g_1^2, \\
16\pi^2\gamma_L^{(1)} &= E - \tfrac{3}{2}g_2^2 - \tfrac{3}{10}g_1^2, \\
16\pi^2\gamma_\tau^{(1)} &= 2\tilde{E} - \tfrac{6}{5}g_1^2.
\end{aligned} \tag{7}$$

and at two loops[6]:

$$\begin{aligned}
(16\pi^2)^2\gamma_t^{(2)} &= -2\tilde{T}^2 - 6(\mathrm{tr}T)\tilde{T} - 2Y_t\,BY_t^\dagger + \left(6g_2^2 - \tfrac{2}{5}g_1^2\right)\tilde{T} \\
&\quad + (\tfrac{8}{15}b_1 + \tfrac{64}{225})g_1^4 + \tfrac{128}{45}g_1^2g_3^2 + (\tfrac{8}{3}b_3 + \tfrac{64}{9})g_3^4, && (8a) \\
(16\pi^2)^2\gamma_b^{(2)} &= -2\tilde{B}^2 - 6(\mathrm{tr}B)\tilde{B} - 2Y_b\,TY_b^\dagger - 2(\mathrm{tr}E)\tilde{B} + \left(6g_2^2 + \tfrac{2}{5}g_1^2\right)\tilde{B} \\
&\quad + (\tfrac{2}{15}b_1 + \tfrac{4}{225})g_1^4 + \tfrac{32}{45}g_1^2g_3^2 + (\tfrac{8}{3}b_3 + \tfrac{64}{9})g_3^4, && (8b) \\
(16\pi^2)^2\gamma_Q^{(2)} &= -2T^2 - 3(\mathrm{tr}T)T - 2B^2 - 3(\mathrm{tr}B)B - (\mathrm{tr}E)B + g_1^2(\tfrac{4}{5}T + \tfrac{2}{5}B) + \tfrac{1}{10}g_1^2g_2^2 \\
&\quad + 8g_3^2g_2^2 + \tfrac{8}{45}g_1^2g_3^2 + (\tfrac{8}{3}b_3 + \tfrac{64}{9})g_3^4 + (\tfrac{3}{2}b_2 + \tfrac{9}{4})g_2^4 + (\tfrac{1}{30}b_1 + \tfrac{1}{900})g_1^4, && (8c) \\
(16\pi^2)^2\gamma_{H_2}^{(2)} &= -9\mathrm{tr}T^2 - 3\mathrm{tr}BT + \left(16g_3^2 + \tfrac{4}{5}g_1^2\right)\mathrm{tr}T + (\tfrac{3}{2}b_2 + \tfrac{9}{4})g_2^4 + \tfrac{9}{10}g_1^2g_2^2 \\
&\quad + (\tfrac{3}{10}b_1 + \tfrac{9}{100})g_1^4, && (8d) \\
(16\pi^2)^2\gamma_{H_1}^{(2)} &= -9\mathrm{tr}B^2 - 3\mathrm{tr}BT - 3(\mathrm{tr}E^2) + \left(16g_3^2 - \tfrac{2}{5}g_1^2\right)\mathrm{tr}B + \tfrac{6}{5}g_1^2\mathrm{tr}E \\
&\quad + (\tfrac{3}{2}b_2 + \tfrac{9}{4})g_2^4 + \tfrac{9}{10}g_1^2g_2^2 + (\tfrac{3}{10}b_1 + \tfrac{9}{100})g_1^4, && (8e) \\
(16\pi^2)^2\gamma_L^{(2)} &= -2E^2 - 3(\mathrm{tr}B)E - (\mathrm{tr}E)E + \tfrac{6}{5}g_1^2E + (\tfrac{3}{2}b_2 + \tfrac{9}{4})g_2^4 \\
&\quad + \tfrac{9}{10}g_1^2g_2^2 + (\tfrac{3}{10}b_1 + \tfrac{9}{100})g_1^4, && (8f) \\
(16\pi^2)^2\gamma_\tau^{(2)} &= -2\tilde{E}^2 - 6(\mathrm{tr}B)\tilde{E} - 2(\mathrm{tr}E)\tilde{E} + \left(6g_2^2 - \tfrac{6}{5}g_1^2\right)\tilde{E} + (\tfrac{6}{5}b_1 + \tfrac{36}{25})g_1^4. && (8g)
\end{aligned}$$

The three loop results are[3]:

$$\begin{aligned}
(16\pi^2)^3\gamma_t^{(3)} &= (2\kappa + 6)\tilde{T}^3 + 36(\mathrm{tr}T^2)\tilde{T} + 6(\mathrm{tr}T)\tilde{T}^2 - 18(\mathrm{tr}T)^2\tilde{T} + 12(\mathrm{tr}TB)\tilde{T} \\
&\quad - 2Y_t\,BTY_t^\dagger - 2Y_t\,TBY_t^\dagger - 6(\mathrm{tr}T)Y_t\,BY_t^\dagger + 12(\mathrm{tr}B)Y_t\,BY_t^\dagger + 6Y_t\,B^2Y_t^\dagger \\
&\quad + 4(\mathrm{tr}E)Y_t\,BY_t^\dagger + \left[\tfrac{64}{3}g_3^2 + (9 - 3\kappa)g_2^2 + \left(\kappa - \tfrac{1}{3}\right)g_1^2\right]\tilde{T}^2 + \Big[(16\kappa - 16)\,g_3^2 \\
&\quad - (9\kappa - 27)g_2^2 + \left(7 + \tfrac{7}{5}\kappa\right)g_1^2\Big](\mathrm{tr}T)\tilde{T} + \left[\tfrac{64}{3}g_3^2 - (3\kappa - 9)g_2^2 + \left(\tfrac{19}{15} + \tfrac{3}{5}\kappa\right)g_1^2\right]Y_t\,BY_t^\dagger \\
&\quad + \Big[\left(\tfrac{304}{3} - \tfrac{272}{9}\kappa - 32n_g\right)g_3^4 + (16\kappa - 88)g_2^2g_3^2 - \left(\tfrac{112}{45}\kappa + \tfrac{8}{15}\right)g_1^2g_3^2 \\
&\quad - \left(\tfrac{3}{2}\kappa + 24n_g - \tfrac{57}{2}\right)g_2^4 - \left(\tfrac{67}{5} - \tfrac{13}{5}\kappa\right)g_1^2g_2^2 - \left(\tfrac{247}{450}\kappa + \tfrac{237}{150} + \tfrac{24}{5}n_g\right)g_1^4\Big]\tilde{T} \\
&\quad - \left(\tfrac{80}{3}g_3^4 + \tfrac{104}{15}g_1^4\right)\mathrm{tr}T - \left(\tfrac{56}{15}g_1^4 + \tfrac{80}{3}g_3^4\right)\mathrm{tr}B - \tfrac{24}{5}g_1^4\mathrm{tr}E \\
&\quad + \left(\tfrac{160}{9}\kappa n_g + \tfrac{32}{3}n_g^2 + \tfrac{368}{9}n_g - \tfrac{3184}{27}\right)g_3^6 - (4\kappa n_g - 20n_g)\,g_2^2g_3^4 \\
&\quad + \left(\tfrac{92}{45}n_g - \tfrac{44}{45}\kappa n_g - \tfrac{448}{45}\right)g_1^2g_3^4 - \left(\tfrac{352}{225}\kappa n_g + \tfrac{1216}{225} - \tfrac{224}{45}n_g\right)g_1^4g_3^2 - \Big[\left(\tfrac{12}{25} + \tfrac{8}{25}n_g\right)\kappa \\
&\quad - \tfrac{8}{5}n_g - \tfrac{12}{5}\Big]g_1^4g_2^2 - \Big[\left(\tfrac{12}{125} + \tfrac{152}{225}n_g\right)\kappa - \tfrac{184}{45}n_g - \tfrac{32}{15}n_g^2 - \tfrac{668}{3375}\Big]g_1^6, && (9a) \\
(16\pi^2)^3\gamma_b^{(3)} &= (2\kappa + 6)\tilde{B}^3 + 36(\mathrm{tr}B^2)\tilde{B} + 6(\mathrm{tr}B)\tilde{B}^2 - 18(\mathrm{tr}B)^2\tilde{B} + 12(\mathrm{tr}BT)\tilde{B}
\end{aligned}$$

$$
\begin{aligned}
&- 2Y_b\,TBY_b^\dagger - 2Y_b\,BTY_b^\dagger - 6(\mathrm{tr}B)Y_b\,TY_b^\dagger + 12(\mathrm{tr}T)Y_b\,TY_b^\dagger + 6Y_b\,T^2Y_b^\dagger \\
&+ 12(\mathrm{tr}E^2)\tilde{B} + 2(\mathrm{tr}E)\tilde{B}^2 - 2(\mathrm{tr}E)Y_b\,TY_b^\dagger - 12(\mathrm{tr}E)(\mathrm{tr}B)\tilde{B} - 2(\mathrm{tr}E)^2\tilde{B} \\
&+ \Big[\tfrac{64}{3}g_3^2 - (3\kappa-9)g_2^2 + (\tfrac{3}{5}\kappa - \tfrac{29}{15})g_1^2\Big]Y_b\,TY_b^\dagger + \Big[\tfrac{64}{3}g_3^2 - (3\kappa-9)g_2^2 \\
&+ \big(\tfrac{1}{5}\kappa - \tfrac{1}{3}\big)\,g_1^2\Big]Y_b\,BY_b^\dagger + \Big[(16\kappa-16)\,g_3^2 - (9\kappa-27)g_2^2 + (7-\kappa)\,g_1^2\Big](\mathrm{tr}B)\tilde{B} \\
&+ \Big[16g_3^2 + (9-3\kappa)g_2^2 + (\kappa-3)g_1^2\Big](\mathrm{tr}E)\tilde{B} + \Big[-\big(\tfrac{272}{9}\kappa - \tfrac{304}{3} + 32n_g\big)\,g_3^4 \\
&+ (16\kappa-88)\,g_2^2g_3^2 + \big(\tfrac{16}{9}\kappa - \tfrac{24}{5}\big)\,g_1^2g_3^2 - \big(\tfrac{3}{2}\kappa + 24n_g - \tfrac{57}{2}\big)\,g_2^4 + \big(\tfrac{7}{5}\kappa - \tfrac{43}{5}\big)\,g_1^2g_2^2 \\
&- \big(\tfrac{49}{30} + \tfrac{16}{5}n_g + \tfrac{7}{450}\kappa\big)\,g_1^4\Big]\tilde{B} - \big(\tfrac{26}{15}g_1^4 + \tfrac{80}{3}g_3^4\big)\,\mathrm{tr}T - \Big[\tfrac{80}{3}g_3^4 + \tfrac{14}{15}g_1^4\Big]\mathrm{tr}B \\
&- \tfrac{6}{5}g_1^4\mathrm{tr}\tilde{E} + \big(\tfrac{160}{9}\kappa n_g + \tfrac{32}{3}n_g^2 + \tfrac{368}{9}n_g - \tfrac{3184}{27}\big)\,g_3^6 - \Big[\big(\tfrac{3}{25} + \tfrac{2}{25}n_g\big)\,\kappa - \tfrac{2}{5}n_g \\
&- \tfrac{3}{5}\Big]g_1^4g_2^2 - \big(\tfrac{112}{45} + \tfrac{44}{45}\kappa n_g - \tfrac{188}{45}n_g\big)\,g_1^2g_3^4 - \big(\tfrac{88}{225}\kappa n_g - \tfrac{56}{45}n_g + \tfrac{112}{225}\big)\,g_1^4g_3^2 \\
&- (4\kappa n_g - 20n_g)g_2^2g_3^4 - \Big[\big(\tfrac{38}{225}n_g + \tfrac{3}{125}\big)\,\kappa - \tfrac{8}{15}n_g^2 - \tfrac{254}{225}n_g - \tfrac{103}{675}\Big]g_1^6,
\end{aligned}
\tag{9b}
$$

$$
\begin{aligned}
(16\pi^2)^3\gamma_Q^{(3)} &= \kappa T^3 + 18(\mathrm{tr}T^2)T + 6(\mathrm{tr}T)T^2 - 9(\mathrm{tr}T)^2T + \kappa B^3 + 18(\mathrm{tr}B^2)B + 6(\mathrm{tr}B)B^2 \\
&- 9(\mathrm{tr}B)^2B + 6(\mathrm{tr}BT)T + 4TBT + 6(\mathrm{tr}TB)B + 4BTB + 6(\mathrm{tr}E^2)B - (\mathrm{tr}E)^2B \\
&+ 2(\mathrm{tr}E)B^2 - 6(\mathrm{tr}B)(\mathrm{tr}E)B + \Big[\tfrac{64}{3}g_3^2 + (3\kappa-3)g_2^2 - \big(\kappa - \tfrac{11}{3}\big)\,g_1^2\Big]T^2 \\
&+ \Big[(8\kappa-8)g_3^2 + 18g_2^2 - \big(\tfrac{4}{5}\kappa - 2\big)\,g_1^2\Big](\mathrm{tr}T)T + \Big[\tfrac{64}{3}g_3^2 + (3\kappa-3)g_2^2 - \big(\tfrac{1}{5}\kappa - \tfrac{7}{15}\big)\,g_1^2\Big]B^2 \\
&+ \Big[(8\kappa-8)g_3^2 + 18g_2^2 - \big(\tfrac{4}{5}\kappa - \tfrac{16}{5}\big)\,g_1^2\Big](\mathrm{tr}B)B + \Big[8g_3^2 + 6g_2^2 + \big(\tfrac{2}{5}\kappa - \tfrac{8}{5}\big)\,g_1^2\Big](\mathrm{tr}E)B \\
&+ \Big[-\big(\tfrac{136}{9}\kappa + 16n_g - \tfrac{152}{3}\big)\,g_3^4 - 4g_2^2g_3^2 - \big(\tfrac{68}{5} - \tfrac{64}{45}\kappa\big)\,g_1^2g_3^2 - \big(9n_g + \tfrac{21}{4}\kappa - \tfrac{63}{4}\big)\,g_2^4 \\
&+ \big(\tfrac{3}{2}\kappa - \tfrac{59}{10}\big)\,g_1^2g_2^2 + \big(\tfrac{143}{900}\kappa - \tfrac{707}{300} - \tfrac{17}{5}n_g\big)\,g_1^4\Big]T + \Big[-\big(\tfrac{136}{9}\kappa + 16n_g - \tfrac{152}{3}\big)\,g_3^4 \\
&- 4g_2^2g_3^2 + \big(\tfrac{64}{45}\kappa - \tfrac{76}{15}\big)\,g_1^2g_3^2 - \big(9n_g + \tfrac{21}{4}\kappa - \tfrac{63}{4}\big)\,g_2^4 + \big(\tfrac{3}{10}\kappa - \tfrac{41}{10}\big)\,g_1^2g_2^2 \\
&- \big(\tfrac{9}{5}n_g + \tfrac{279}{300} - \tfrac{7}{180}\kappa\big)\,g_1^4\Big]B - \Big[\tfrac{80}{3}g_3^4 + \tfrac{45}{2}g_2^4 + \tfrac{13}{30}g_1^4\Big]\mathrm{tr}T \\
&- \Big[\tfrac{80}{3}g_3^4 + \tfrac{45}{2}g_2^4 + \tfrac{7}{30}g_1^4\Big]\mathrm{tr}B - \big(\tfrac{15}{2}g_2^4 + \tfrac{3}{10}g_1^4\big)\,\mathrm{tr}E + \Big[\tfrac{160}{9}n_g\kappa + \tfrac{32}{3}n_g^2 \\
&+ \tfrac{368}{9}n_g - \tfrac{3184}{27}\Big]g_3^6 + (12n_g - 4\kappa n_g - 28)\,g_2^2g_3^4 + \big(\tfrac{212}{45}n_g - \tfrac{44}{45}\kappa n_g - \tfrac{28}{45}\big)\,g_1^2g_3^4 \\
&- \Big[\big(\tfrac{3}{100} + \tfrac{1}{50}n_g\big)\,\kappa - \tfrac{11}{100}\Big]g_1^4g_2^2 - \tfrac{8}{5}g_1^2g_2^2g_3^2 + \big(\tfrac{14}{45}n_g - \tfrac{22}{225}\kappa n_g - \tfrac{16}{225}\big)\,g_1^4g_3^2 \\
&+ \Big[\big(\tfrac{15}{2}n_g + \tfrac{15}{4}\big)\,\kappa - \tfrac{87}{4} + 18n_g + 6n_g^2\Big]g_2^6 - \Big[\big(\tfrac{9}{20} + \tfrac{3}{10}n_g\big)\,\kappa - \tfrac{7}{5}n_g - \tfrac{41}{20}\Big]g_1^2g_2^4 \\
&- (6\kappa n_g - 22n_g + 16)\,g_2^4g_3^2 - \Big[\big(\tfrac{3}{500} + \tfrac{19}{450}n_g\big)\,\kappa - \tfrac{2}{15}n_g^2 - \tfrac{13}{45}n_g - \tfrac{557}{13500}\Big]g_1^6,
\end{aligned}
\tag{9c}
$$

$$
\begin{aligned}
(16\pi^2)^3\gamma_{H_2}^{(3)} &= (3\kappa+3)\mathrm{tr}T^3 + 54\mathrm{tr}T\mathrm{tr}T^2 + 9\mathrm{tr}TB^2 + 18\mathrm{tr}B\mathrm{tr}BT + 6\mathrm{tr}E\mathrm{tr}BT \\
&+ \Big[(72-24\kappa)g_3^2 + (9+9\kappa)g_2^2 + \big(\tfrac{57}{5} - \tfrac{3}{5}\kappa\big)\,g_1^2\Big]\mathrm{tr}T^2 + \Big[(24-8\kappa)g_3^2 + 18g_2^2
\end{aligned}
$$

$$
\begin{aligned}
&+\big(\tfrac{1}{5}\kappa+\tfrac{6}{5}\big)\,g_1^2\Big]\mathrm{tr}BT-\Big[\big(64n_g+\tfrac{8}{3}\kappa-\tfrac{416}{3}\big)\,g_3^4+\big(27n_g-\tfrac{99}{4}+\tfrac{63}{4}\kappa\big)\,g_2^4\\
&+\big(\tfrac{43}{5}n_g+\tfrac{115}{12}+\tfrac{13}{60}\kappa\big)\,g_1^4-(24\kappa-132)\,g_2^2g_3^2-\big(\tfrac{104}{15}\kappa-\tfrac{124}{3}\big)\,g_1^2g_3^2\\
&-\big(\tfrac{21}{10}\kappa-\tfrac{57}{10}\big)\,g_1^2g_2^2\Big]\mathrm{tr}T-\big(\tfrac{45}{2}g_2^4+\tfrac{21}{10}g_1^4\big)\,\mathrm{tr}B-\big(\tfrac{15}{2}g_2^4+\tfrac{27}{10}g_1^4\big)\,\mathrm{tr}E+\Xi, \qquad (9d)
\end{aligned}
$$

$$
\begin{aligned}
(16\pi^2)^3\gamma_{H_1}^{(3)}&=(3\kappa+3)\mathrm{tr}B^3+54\mathrm{tr}B\mathrm{tr}B^2+18\mathrm{tr}T\mathrm{tr}BT+9\mathrm{tr}BT^2+(\kappa+1)\mathrm{tr}E^3\\
&+6\mathrm{tr}E\mathrm{tr}E^2+18\mathrm{tr}B\mathrm{tr}E^2+18\mathrm{tr}E\mathrm{tr}B^2+\Big[\big(\tfrac{7}{5}\kappa-\tfrac{12}{5}\big)\,g_1^2\\
&+(24-8\kappa)g_3^2+18g_2^2\Big]\mathrm{tr}BT+\Big[(72-24\kappa)g_3^2+(9+9\kappa)g_2^2+\big(\tfrac{9}{5}\kappa+3\big)\,g_1^2\Big]\mathrm{tr}B^2\\
&+\Big[(3\kappa+3)\,g_2^2-\big(\tfrac{9}{5}\kappa-9\big)\,g_1^2\Big](\mathrm{tr}E^2)-\big(\tfrac{45}{2}g_2^4+\tfrac{39}{10}g_1^4\big)\,\mathrm{tr}T+\Big[\big(\tfrac{416}{3}-64n_g-\tfrac{8}{3}\kappa\big)\,g_3^4\\
&+(24\kappa-132)\,g_2^2g_3^2+\big(\tfrac{56}{15}\kappa-\tfrac{284}{15}\big)\,g_1^2g_3^2-\big(\tfrac{63}{4}\kappa-\tfrac{99}{4}+27n_g\big)\,g_2^4\\
&-\big(\tfrac{3}{2}\kappa+\tfrac{3}{10}\big)\,g_1^2g_2^2-\big(\tfrac{77}{300}\kappa+\tfrac{191}{60}+\tfrac{19}{5}n_g\big)\,g_1^4\Big]\mathrm{tr}B+\Big[\big(\tfrac{33}{4}-\tfrac{21}{4}\kappa-9n_g\big)\,g_2^4\\
&+\big(\tfrac{27}{10}\kappa-\tfrac{81}{10}\big)\,g_1^2g_2^2+\big(\tfrac{27}{100}\kappa-\tfrac{33}{5}n_g-\tfrac{207}{20}\big)\,g_1^4\Big]\mathrm{tr}E+\Xi, \qquad (9e)
\end{aligned}
$$

$$
\begin{aligned}
(16\pi^2)^3\gamma_{L}^{(3)}&=\kappa E^3+6(\mathrm{tr}E^2)E+2(\mathrm{tr}E)E^2-(\mathrm{tr}E)^2E+6(\mathrm{tr}B)E^2-6(\mathrm{tr}B)(\mathrm{tr}E)E\\
&+18(\mathrm{tr}B^2)E+6(\mathrm{tr}BT)E-9(\mathrm{tr}B)^2E+\Big[(3\kappa-3)g_2^2-\big(\tfrac{9}{5}\kappa-9\big)\,g_1^2\Big]E^2\\
&+\Big[(8\kappa-32)g_3^2+18g_2^2-(2\kappa-8)g_1^2\Big](\mathrm{tr}B)E+6g_2^2(\mathrm{tr}E)E-\big(\tfrac{39}{10}g_1^4+\tfrac{45}{2}g_2^4\big)\,\mathrm{tr}T\\
&+\Big[\big(\tfrac{63}{4}-\tfrac{21}{4}\kappa-9n_g\big)\,g_2^4+\big(\tfrac{27}{10}\kappa-\tfrac{81}{10}\big)\,g_1^2g_2^2+\big(\tfrac{27}{100}\kappa-\tfrac{153}{20}-\tfrac{33}{5}n_g\big)\,g_1^4\Big]E\\
&-\Big[\tfrac{21}{10}g_1^4+\tfrac{45}{2}g_2^4\Big]\mathrm{tr}B-\Big[\tfrac{15}{2}g_2^4+\tfrac{27}{10}g_1^4\Big]\mathrm{tr}E+\Xi, \qquad (9f)
\end{aligned}
$$

$$
\begin{aligned}
(16\pi^2)^3\gamma_{\tau}^{(3)}&=(2\kappa+6)\tilde{E}^3+12(\mathrm{tr}E^2)\tilde{E}+2(\mathrm{tr}E)\tilde{E}^2-2(\mathrm{tr}E)^2\tilde{E}+6(\mathrm{tr}B)\tilde{E}^2\\
&-12(\mathrm{tr}B)(\mathrm{tr}E)\tilde{E}+36(\mathrm{tr}B^2)\tilde{E}-18(\mathrm{tr}B)^2\tilde{E}+12(\mathrm{tr}BT)\tilde{E}+\Big[(9-3\kappa)g_2^2\\
&+(\tfrac{9}{5}\kappa+\tfrac{9}{5})g_1^2\Big]\Big[\tilde{E}^2+(\mathrm{tr}E)\tilde{E}\Big]+\Big[(16\kappa-64)\,g_3^2-(9\kappa-27)\,g_2^2\\
&+\big(\tfrac{107}{5}+\tfrac{7}{5}\kappa\big)\,g_1^2\Big](\mathrm{tr}B)\tilde{E}-\tfrac{78}{5}g_1^4\mathrm{tr}T-\tfrac{42}{5}g_1^4\mathrm{tr}B-\tfrac{54}{5}g_1^4\mathrm{tr}E\\
&+\Big[-\big(\tfrac{3}{2}\kappa+24n_g-\tfrac{57}{2}\big)\,g_2^4+\big(\tfrac{27}{5}\kappa-27\big)\,g_1^2g_2^2-\big(\tfrac{63}{50}+\tfrac{27}{10}\kappa+\tfrac{48}{5}n_g\big)\,g_1^4\Big]\tilde{E}\\
&+\big(\tfrac{88}{5}n_g-\tfrac{88}{25}\kappa n_g\big)\,g_1^4g_3^2+\Big[\tfrac{18}{5}n_g+\tfrac{27}{5}-\big(\tfrac{27}{25}+\tfrac{18}{25}n_g\big)\,\kappa\Big]g_1^4g_2^2\\
&+\Big[\tfrac{24}{5}n_g^2+\tfrac{38}{5}n_g-\big(\tfrac{38}{25}n_g+\tfrac{27}{125}\big)\,\kappa-\tfrac{351}{125}\Big]g_1^6, \qquad (9g)
\end{aligned}
$$

where $\kappa=6\zeta(3)$, and

$$
\begin{aligned}
\Xi&=n_g\,(30-6\kappa)\,g_2^4g_3^2+n_g\,\big(\tfrac{22}{5}-\tfrac{22}{25}\kappa\big)\,g_1^4g_3^2+\Big[\big(\tfrac{15}{2}n_g+\tfrac{15}{4}\big)\,\kappa-\tfrac{87}{4}+18n_g+6n_g^2\Big]g_2^6\\
&-\Big[\big(\tfrac{9}{20}+\tfrac{3}{10}n_g\big)\,\kappa-\tfrac{9}{20}-\tfrac{3}{5}n_g\Big]g_1^2g_2^4-\Big[\big(\tfrac{9}{50}n_g+\tfrac{27}{100}\big)\,\kappa-\tfrac{27}{100}\Big]g_1^4g_2^2\\
&-\Big[\big(\tfrac{19}{50}n_g+\tfrac{27}{500}\big)\,\kappa-\tfrac{27}{100}-\tfrac{6}{5}n_g^2-\tfrac{61}{25}n_g\Big]g_1^6. \qquad (10)
\end{aligned}
$$

In terms of the anomalous dimensions, the Yukawa β-functions are:

$$\beta_{Y_t} = \gamma_t Y_t + Y_t(\gamma_Q + \gamma_{H_2}), \quad \beta_{Y_b} = \gamma_b Y_b + Y_b(\gamma_Q + \gamma_{H_1}), \quad \beta_{Y_\tau} = \gamma_\tau Y_\tau + Y_\tau(\gamma_L + \gamma_{H_1}). \tag{11}$$

In the approximation that we retain only α_3 and the t-quark Yukawa coupling y_t then we have (for $n_g = 3$):

$$16\pi^2 \beta^{(1)}_{y_t} = y_t(6y_t^2 - \tfrac{16}{3}g_3^2), \tag{12a}$$

$$(16\pi^2)^2 \beta^{(2)}_{y_t} = y_t(-22y_t^4 + 16y_t^2 g_3^2 - \tfrac{16}{9}g_3^4), \tag{12b}$$

$$(16\pi^2)^3 \beta^{(3)}_{y_t} = y_t([102 + 6\kappa]y_t^6 + \tfrac{272}{3}y_t^4 g_3^2 - [\tfrac{296}{3} + 48\kappa]y_t^2 g_3^4 + [\tfrac{5440}{27} + \tfrac{320}{3}\kappa]g_3^6). \tag{12c}$$

We see that for values of y_t in the neighbourhood of or greater than the one-loop QFP (which corresponds to $y_t \approx 1$) we have $\beta^{(2)}_{y_t} < 0$ and $\beta^{(3)}_{y_t} > 0$. We may therefore expect that where three loop contributions are not completely negligible they will tend to cancel the two loop contributions. We will see examples of this behaviour presently.



Fig.1: Plot of $y_t(M_G)$ against $y_t(M_Z)$. The solid, dashed and dotted lines correspond to one, two and three-loop β-functions respectively.

For small $\tan\beta$, QFP behaviour is of interest because there is a large domain of possible values of $y_t(M_G)$ that lead to the same value of $y_t(M_Z)$. (For a recent discussion, see Ref. [7].) It is important, however, to consider to what extent this domain is restricted by the requirement of perturbative believability. Setting $g_3 = 0$ above, we see that $|\beta^{(2)}_{Y_t}/\beta^{(1)}_{Y_t}| \approx 1$ for $y_t \approx 6.6$ while $|\beta^{(3)}_{Y_t}/\beta^{(2)}_{Y_t}| \approx 1$ for $y_t \approx 4.9$. So since the QFP corresponds to $y_t \approx 1.1$, we may expect there to be a good sized domain available for $y_t(M_G)$ such that we have both perturbative believability and approach to the QFP at low energies. This is illustrated in Figure 1, where (with $m_t^{\rm pole} = 175$GeV, and using the complete β-functions, not the Eq. (12) approximations) we plot $y_t(M_G)$ against $y_t(M_Z)$, for values close to the QFP. The breakdown of perturbation theory at $y_t(M_G) \approx 6$ is clearly seen.

In our running analysis † we take the effective field theory to be the SSM for all scales between M_Z and M_G: see Ref. [9] for a discussion of this procedure. A reason to prefer this to the traditional stepwise method is that the latter involves non-supersymmetric intermediate theories for which (beyond one loop) use of DRED rather than DREG is problematic [10]. This choice, means, of course, that the input values of $\alpha_{1\to 3}$ are sensitive to our assumptions about the sparticle spectrum. We have corrected for this at one-loop using Ref. [9]; of course we should use two-loop corrections for consistency, as we should, for example, in converting the running m_t to the pole m_t. These calculations remain to be done however; there exists a result for the two-loop gluon contribution to $m_t^{\rm pole}$ but this is in DREG not DRED. We do not impose trinification of the gauge couplings, since this leads to somewhat high values of $\alpha_3(M_Z)$; for a discussion and references see Ref. [11]. We input α_3 and define unification to be where α_1 and α_2 meet. Our input values at M_Z are $\alpha_1 = 0.0167$, $\alpha_2 = 0.032$ and $\alpha_3 = 0.1$. These values correspond to a superpartner spectrum with an effective supersymmetric scale[12]of $T_{\rm SUSY} = 1$TeV, and $\alpha_3^{\rm SM} = 0.117$.



Fig.2: Plot of $\tan\beta$ *against* $m_t^{\rm pole}$ *with Yukawa trinification. The solid, dashed and dotted lines correspond to one, two and three-loop* β*-functions respectively.*

---

† For a different approach to this running analysis based on the NSVZ β_g, see Ref. [8].

In Figure 2 we plot $\tan\beta$ against m_t assuming trinification ($y_t = y_b = y_\tau$) at M_G. As anticipated, the three loop corrections counteract (to some extent) the two loop ones. Their effect, while small, is not completely negligible. Proximity of y_t and y_b to their QFPs leads to an upper limit $m_t \approx 181$GeV. (In comparing with, for example Fig. 1 of Ref. [13], it is important to note that there they have taken the sparticle masses to be at M_Z.)



Fig.3: Plot of $m_b^{\rm pole}$ *against* $\tan\beta$. *The solid, dashed and dotted lines correspond to one, two and three-loop* β*-functions respectively.*

As another example, consider the low $\tan\beta$ region. In Figure 3 we plot $m_b^{\rm pole}$ against $\tan\beta$, where we have imposed $b-\tau$ unification. (For running from M_Z to m_b we use the $SU_3 \otimes U_1$ β-functions given, for example in Ref. [14].) In this graph the QFP is approached as $\tan\beta$ decreases. (The fact that proximity to the QFP gives a better value of $m_b^{\rm pole}$ has been noted by a number of authors; for an early example see Ref. [6].) The result for $m_b^{\rm pole}$ is quite sensitive to the input value of α_3 (and hence to the sparticle spectrum).

In conclusion: detailed running coupling analyses for the dimensionless SSM couplings have been performed by a number of groups; we have not here duplicated in full these efforts, but instead investigated the effect on them of the three loop β-functions. We have seen that the corrections are small; nevertheless it is possible that one day they will play a part in a very accurate comparison between the SSM and experiment.

In non-minimal models, the three-loop terms may assume more immediate importance. Specifically, consider the possibility that $\beta_{g_3}^{(1)} = 0$. In such cases two-loop corrections will clearly dominate the evolution of g_3. It was shown in Ref. [6] that at two loops perturbative unification is only just achievable. Evidently three-loop contributions will be important here, and in fact improve matters. Models of this kind will have interesting phenomenology, with significant differences from the SSM.[15]

Acknowledgements

IJ was supported by PPARC via an Advanced Fellowship, and PF by a scholarship from JNICT. We thank Chris Kolda, John March-Russell and Damien Pierce for conversations.

References

[1] B. Pendleton and G.G. Ross, Phys. Lett. **B**98 (1981) 291;
C.T. Hill, Phys. Rev. **D**24 (1981) 691

[2] B. Ananthanarayan, G. Lazarides and Q. Shafi, Phys. Rev. **D**44 (1991) 1613;
H. Arason et al, Phys. Rev. Lett. 67 (1991) 2933;
V. Barger, M.S. Berger and P. Ohmann, Phys. Rev. **D**47 (1993) 1093;
M. Carena et al, Nucl. Phys. **B**426 (1994) 269;
L.J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. **D**50 (1994) 7048;
R. Rattazzi and U. Sarid, Phys. Rev. **D**53 (1996) 1553;
E.G. Floratos, G.K. Leontaris and S. Lola, Phys. Lett. **B**365 (1996) 149

[3] I. Jack, D.R.T. Jones and C.G. North, hep-ph/9603386

[4] I. Jack, D.R.T. Jones and C.G. North, hep-ph/9606323

[5] V.A. Novikov et al, Phys. Lett. **B**166 (1986) 329;
M.A. Shifman and A.I Vainstein, Nucl. Phys. **B**277 (1986) 456

[6] J.E. Björkman and D.R.T. Jones, Nucl. Phys. **B**259 (1985) 533

[7] M. Lanzagorta and G.G. Ross, Phys. Lett. **B**349 (1995) 319

[8] M. Shifman, hep-ph/9606281

[9] A.E. Faraggi and B. Grinstein, Nucl. Phys. **B**422 (1994) 3;
P.H. Chankowski, Z. Pluciennik and S. Pokorski, Nucl. Phys. **B**439 (1995) 23;
J. Bagger, K. Matchev and D. Pierce, Phys. Lett. **B**348 (1995) 443;
D. Pierce et al, hep-ph/9606211

[10] I. Jack, D.R.T. Jones and K.L. Roberts, Z. Phys. **C**62 (1994) 161; *ibid***C**63 (1994) 151

[11] L. Roszkowski, hep-ph/9509273

[12] P. Langacker and N. Polonski, Phys. Rev. **D**47 (1993) 4028;
M. Carena, S. Pokorski and C.E.M. Wagner, Nucl. Phys. **B**406 (1993) 59

[13] M. Carena et al, Ref. [2]

[14] H. Arason et al, Phys. Rev. **D**46 (1992) 3945

[15] C. Kolda and J. March-Russell, to be published